§ 52.1853

additional requirements as outlined in Table II of this subpart.

(c) "U.S. Grade C" is the quality of Raisins with Seeds that have similar varietal characteristics; that have a fairly good typical color with not more than 20 percent, by weight, that may be dark reddish-brown berries; that have a fairly good flavor; that show development characteristics of raisins prepared from fairly well-matured grapes;

that contain not more than 18 percent, by weight, of moisture, except that any seeded raisins may contain not more than 19 percent, by weight, of moisture; and meet the additional requirements as outlined in Table II of this subpart.

(d) "Substandard" is the quality of Raisins with Seeds that fail to meet the requirements of U.S. Grade C.

TABLE II—ALLOWANCES FOR DEFECTS IN RAISINS WITH SEEDS—EXCEPT LAYER OR CLUSTER

| Defects | U.S. Grade A | U.S. Grade B | U.S. Grade C | | | |
|---|--|---|---|--|--|--|
| | Maximum count (per 32 ounces) | | | | | |
| Pieces of stem | 7 | 2 | 3 | | | |
| | Maximum count (per 16 ounces) | | | | | |
| Capstems in other than | | | | | | |
| uncapstemmed types | 10 | 15 | 20 | | | |
| Seeds in seeded types Loose capstems in uncapstemmed | 12 | 15 | 20 | | | |
| types | 20 | 20 | 20 | | | |
| | Maximum (percent by weight) | | | | | |
| Sugared | 5 | 10 | 15 | | | |
| Discolored, damaged, or moldy Provided these limits are not exceeded: | 5 | 7 | 9 | | | |
| Damaged | 3 | 4 | 5 | | | |
| Moldy | 2 | 3 | 4 | | | |
| Substandard development and un- | • | _ | | | | |
| developed | 2 | 5 | 8 | | | |
| | Appearance or edibility of product | | | | | |
| Slightly discolored or damaged by fermentation or any other defect not described above. | May not be affected | May not be more than slightly affected. | May not be more than materially affected. | | | |
| Grit, sand, or silt | None of any consequence m appearance or edibility of the | Not more than a trace ma be present that affects the appearance or edibility of the product. | | | | |

§ 52.1853 Grades of raisins with seeds—layer or cluster.

(a) "U.S. Grade A" is the quality of Layer or Cluster Raisins with Seeds that have similar varietal characteristics; that have a good typical color; that have a good characteristic flavor; that are uniformly cured and show development characteristics of raisins prepared from well-matured grapes; that contain not more than 23 percent, by weight, of moisture; that not less

than 30 percent, by weight, of the raisins, exclusive of stems and branches, are 3-Crown size or larger; and that meet the following additional requirements as outlined in Table III of this subpart.

(b) "U.S. Grade B" is the quality of Layer or Cluster Raisins with Seeds that have similar varietal characteristics that have a reasonably good typical color; that have a good characteristic flavor; that are uniformly cured and

show development characteristics of raisins prepared from reasonably wellmatured grapes; that contain not more than 23 percent, by weight, of the moisture, that not less than 30 percent, by weight, of the raisins exclusive of stems and branches, are 3-Crown size or larger; and that meet the additional requirements outlined in Table III of this subpart.

(c) "Substandard" is the quality of Layer or Cluster Raisins with Seeds that fail to meet the requirements of U.S. Grade B.

TABLE III—ALLOWANCE FOR DEFECTS IN LAYER OR CLUSTER RAISINS WITH SEEDS

| Defects | U.S. Grade A | U.S. Grade B | |
|---|---|--------------|--|
| | Maximum (percent by weight) | | |
| Sugared | 5 | 10 | |
| Discolored, damaged, or moldy | 5 | 7 | |
| Damaged | 3 | 4 | |
| Moldy | 2 | 3 | |
| Substandard development and undeveloped | 2 | 5 | |
| Shattered (or loose) individual berries and small clusters of 2 or 3 berries each. | Practically free Reasonably free. Appearance or edibility of product | | |
| | | | |
| Slightly discolored or damaged by fermentation or any other defect not described above. | May not be affected May not be more than slightly affected. | | |
| Grit, sand, or silt | None of any consequence m appearance or edibility of the | | |

TYPE IV—SULTANA RAISINS

§52.1854 Sizes of Sultana raisins.

Size designations are not applicable to Sultana Raisins.

§52.1855 Grades of Sultana raisins.

(a) "U.S. Grade A" is the quality of Sultana Raisins that have similar varietal characteristics; that have a good typical color; that have a good characteristic flavor; that show development characteristics of raisins prepared from well-matured grapes; with not less than 80 percent, by weight, of raisins that are well-matured or reasonably well-matured; and that contain not more than 18 percent, by weight, of

moisture, and that meet the additional requirements outlined in Table IV of this subpart.

(b) "U.S. Grade B" is the quality of Sultana Raisins that have similar varietal characteristics; that have a reasonably good typical color; that have a good characteristic flavor; that show development characteristics of raisins prepared from reasonably well-matured grapes; with not less than 70 percent, by weight, of raisins that are well-matured or reasonably well-matured; and that contain not more than 18 percent, by weight, of moisture, and that meet the additional requirements as outlined in Table IV of this subpart.

TABLE IV-ALLOWANCES FOR DEFECTS IN SULTANA RAISINS

| Defects | U.S. Grade A | U.S. Grade B | Grade B U.S. Grade C | | |
|----------------|-------------------------------|--------------|----------------------|---|--|
| | Maximum count (per 32 ounces) | | | | |
| Pieces of stem | 1 | | 2 | 3 | |